

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/389135874>

HEIX1 Mutation Effects on Endoplasmic Reticulum Stress, Caspase Activation, and JNK2 Pathways

Article in Journal of Babol University of Medical Sciences · January 2025

CITATIONS

0

READS

20

5 authors, including:



Maytham Abdulkadhim Dragh
University of Misan

34 PUBLICATIONS 422 CITATIONS

SEE PROFILE



Zainab S. Al-Allak
University of Misan

7 PUBLICATIONS 26 CITATIONS

SEE PROFILE



Zainab Zamil Gataa Allami
University of Misan

10 PUBLICATIONS 15 CITATIONS

SEE PROFILE



HEIX1 Mutation Effects on Endoplasmic Reticulum Stress, Caspase Activation, and JNK2 Pathways

M. A. Dragh (PhD)^{*1}, Z. S. Al-Allak (PhD)², Z. Z. G. Allami (MSc)¹

1. Department of Biology, College of Science, University of Misan, Misan, Iraq.

2. College of Dentistry, University of Misan, Misan, Iraq

*Corresponding Author: M. A. Dragh (PhD)

Address: Department of Biology, College of Science, University of Misan, Misan, Iraq.

Tel: +964 (773) 7232342. E-mail: maithamdragh@uomisan.edu.iq

Article Type

ABSTRACT

Research Paper

Background and Objective: Schneider 2 (S2) cells, derived from *Drosophila melanogaster*, are extensively utilized in developmental biology and genetics engineering research. Proper tissue formation depends on the regulation of developmental signalling, with the unfolded protein response (UPR) and autophagy playing critical roles in maintaining endoplasmic reticulum (ER) and mitochondrial homeostasis. Mutations in the HEIX1 gene disrupt these processes, triggering activation of the P-ERK and JNK signalling pathways, which lead to ER stress, mitochondrial dysfunction, and apoptosis. This study examines the molecular mechanisms underlying HEIX1 loss-of-function mutations in S2 cells, focusing on P-ERK and JNK pathway activation and their effects on cellular stress responses.

Methods: This experimental study includes four groups of S2 cells, with 10 samples per group: (1) wild-type (WT) control, (2) HEIX1 homozygous mutant (HEIX1^{-/-}), (3) HEIX1 heterozygous rescue group (*heix1/Df; UAS, Gal4, tubp>HEIX1), and (4) a group treated with the P-ERK inhibitor GSK. Protein extraction and analysis were performed across all groups. Variables such as P-ERK and JNK activation, reactive oxygen species (ROS) levels, apoptosis markers (caspase 9 activation), and UPR-related gene expression (GRP78 and CHOP) were measured using western blotting, immunofluorescence, and ROS assays.

Findings: Loss of HEIX1 function significantly activated the P-ERK pathway, as evidenced by increased P-ERK phosphorylation, caspase 9 activation, and apoptosis. Mutants showed disrupted ER and mitochondrial homeostasis, including swelling and oxidative stress. Rescuing HEIX1 restored normal signalling and reduced apoptosis. P-ERK inhibition accelerated apoptosis and suppressed UPR-related gene expression, underscoring HEIX1's role in proteostasis.

Conclusion: According to the results of this study, the HEIX1 gene is essential for maintaining ER and mitochondrial homeostasis, regulating stress responses, and preventing apoptosis. Its loss leads to P-ERK pathway activation, ER stress, and cell death. These findings provide insights into *Drosophila* development and the broader implications of HEIX1 in understanding human diseases linked to ER stress and apoptosis.

Keywords: Mitochondria, HEIX1 Mutations, ERK Signalling Pathway, Apoptosis-Induced Death, Written by JNK.

Received:

Sep 16th 2024


Revised:

Oct 23rd 2024

Accepted:

Nov 23rd 2024

Cite this article: Dragh MA, Al-Allak ZS, Allami ZZG. HEIX1 Mutation Effects on Endoplasmic Reticulum Stress, Caspase Activation, and JNK2 Pathways. *Journal of Babol University of Medical Sciences*. 2025; 27: e.



Introduction

Schneider 2 (S2) received from *Drosophila Melanogaster* has emerged as unavoidable tools in developmental biology, genetics, medical, disease modeling and stem cell research. Late-stage d. These cells obtained from the melanogaster fetus are a widely accepted model system due to the small size of the fruit fly, small life cycle, high fertility rate and ease of genetic manipulation. In multicellular organisms, *d. Melanogaster* is specifically capable of enabling phenotypic screening for mutation affecting cell behavior in any developmental state, making it an important platform to clarify gene function and regulation (1-3).

The formation of tissues and organs depends on accurately regulated developmental signaling routes. These routes control important cellular processes like development, division and discrimination. Research By taking advantage of its genetic in *Drosophila*, our understanding of these processes has become quite advanced. Manipulations. Techniques such as P-Elements Insertion and Chemical Mutation like Ethylmethyl Sulfonate (EMS) has enabled the generation of targeted mutations, facilitates detailed studies Gene function. While P-elements provide direct mutation, chemical mutation like EMS Offer high efficiency and low bias, which gives them ideal equipment for major developmental checks Jean (4, 5).

An important aspect of cellular homeostasis includes unfolded protein response (UPR) and autophagy, Which are necessary to maintain endoplasmic reticulum (ER) and mitochondrial function. UPR is ER active in response to stress, which occurs when the protein fold within the ER is interrupted. it Pathway restores cellular function by repairing the misfolded protein or, in cases of prolonged stress, motivates Apoptosis to eliminate damaged cells (6). ER plays an important role in protein synthesis, folding, and in Trafficking, while mitochondria provides the necessary energy for these processes. Interaction between ER and mitochondria are facilitated by mitochondrial-jocked membrane (MAMs), special structures It coordinates calcium transfer, lipid metabolism and protein homeostasis. Appropriate er-mitochondrial Communication is necessary to maintain cellular proteostasis and survival (7-9).

Interruption of ER or mitochondrial function can activate the UPR signaling route. These include three major branches: P-ERK, Ire1 α and ATF6. Each branch controls gene expression through separate transcription factors, helping the cell to adapt to stress. However, highly Reactive Oxygen Species (ROS) production can increase stress of ER and give rise to Apoptosis. Protein such as Redox signaling mediator disulfide isomerase (PDI) and (ER) oxidation (ERO1) play a required role in this process, as well as low and oxidation glutathione (GSH/GSSG), NAD (P) H Oxidase 4 (NOX 4), and Calnexin, Along with, Voltage-dependent Pore (VDAC) (10,11). Studies during stress reactions have highlighted the role of Mitochondrial UPR (UPRMT) in a newly identified route in conversation between mitochondria and nucleus (10).

The P-ERK signaling route is an important component of UPR and is particularly active during stress conditions such as hypoxia and nutrient deficiency. This route is also implicated in tumor metabolism and cellular homeostasis. Tere1 genes in humans and its *Drosophila* Homolog, heix1 have provided valuable insight into the roles of these routes. Tere1 is added to cholesterol metabolism and redox cycling of vitamin K, while the heix1 is shown to regulate the ER-Mitochondrial function. Loss-of-function mutations in Tere1 are associated with diseases such as bladder and prostate cancer, while heix1 mutations increase ATP production in hemocyte overproofing, mitochondrial dysfunction and *drosophila* (12–16). Given its high sequence similarity to the human tere1, the heix1 provides a model to study the ER-Mitochondrial Interaction and their implications in the disease. The heix1 inserts mutations of mutation ER and mitochondrial homeostasis, causing ER stress, activation of P-ERK and JNK signaling routes, and apoptosis (17).

These findings suggest an important role for the *heix1* in maintaining protrustesis and regulating cellular stress reactions. The purpose of this study is to examine the activation of P-ERK and JNK signaling routes in response to the *heix1* loss-of-function mutation in 2 cells. In particular, it examines the regulatory system of developmental signaling, UPR in ER and mitochondria, and ER stress and interconnection between tension and mitochondrial function. By clarifying these mechanisms, the study wants to provide insight into the role of cell behavior, development, and apoptosis, with a wide implication for developmental biology, genetics and disease research.

Methods

This experimental study was conducted in Genetic Engineering Laboratory, Department of Biology, The College of Science, University of Misan, Iraq between September 2023 and April 2024. Including sample size and statistical analysis, was developed in consultation with a statistician to ensure that Working accuracy. Misan Ethics Committee University was obtained from the university (Ethical Code #DepT362), and research followed the principles of the announcement of Helsinki. Written Informed consent was safe from all participants, and strict adherence to moral standards, including To avoid literary theft, data construction and duplicate publication, the entire time was maintained.

Schneider 2 (S2) cells were used as the experimental model, with four distinct groups included in the study. The wild-type (WT) control group was derived from Bloomington stock 1. The *heix1* homozygous mutant group (*HEIX1*^{-/-}) utilized two alleles: a P-element allele from Bloomington stock 11031 and an EMS allele from Bloomington stock 3600. The rescue group involved **heix1*/Df; UAS, Gal4, *tubp*>*HEIX1*, obtained from Bloomington stock 6915. The treatment group consisted of *HEIX1*^{-/-} cells treated with the P-ERK inhibitor Glycogen Synthase Kinase (GSK). The groups were matched to ensure comparability of conditions. Variables examined included P-ERK and JNK pathway activation, reactive oxygen species (ROS) levels, apoptosis markers (caspase 9 activation), and unfolded protein response (UPR)-related gene expression (GRP78 and CHOP).

Total protein was extracted from S2 cells using PROPREP protein extraction buffer (iNTRON Biotechnology, Korea). Samples were centrifuged at 10,000 g for 30 minutes at 4°C, and protein concentrations were determined using a Nanodrop spectrophotometer (Thermo Science, USA). Proteins were prepared in SDS-PAGE loading buffer, separated in a 12% SDS-polyacrylamide gel, and transferred onto polyvinylidene fluoride (PVDF) membranes. Membranes were blocked in 5% non-fat dry milk in TBST and incubated overnight at 4°C with primary antibodies, including Phospho-p44/42 MAPK (P-ERK), p44/42 MAPK (ERK), and Rabbit Polyclonal Caspase-9 antibodies (Abcam). ROS levels were measured using the ROS Assay Kit (Abcam) and H2DCFDA to assess oxidative stress.

Secondary antibodies (Anti-rabbit HRP) were used for immunodetection, followed by washing and development with SuperSignal substrate (Applied Biosystem). Protein bands were visualized on X-ray films and analyzed using Quantitative One Image Software for densitometric quantification. Key measurements included the activation status of P-ERK and ERK, which are critical indicators of UPR involvement (18).

Statistical analysis was conducted with the guidance of a statistician. Data were analyzed using appropriate statistical tests, with a significance level of $p < 0.05$. The choice of tests depended on data normality and distribution, ensuring the accuracy of comparisons between groups and correlations among variables.

Results

The analysis revealed significant differences in P-ERK, total ERK, HEIX1, and ATP5s levels across wild-type (WT), *heix1/heix1* mutants, and rescue groups (**heix1/Df*; UAS, Gal4, *tubp>HEIX1*). In *heix1/heix1* mutants, P-ERK levels were significantly elevated compared to WT (2.7-fold increase, $p=0.002$), indicating hyperactivation of the stress-response pathway. This elevation was reduced to near-WT levels in the rescue group ($p=0.052$), confirming that *heix1* restoration mitigates stress signaling. Total ERK levels remained consistent across all groups, suggesting that the observed changes were specific to P-ERK activation and not due to total ERK protein expression. HEIX1 expression was absent in *heix1/heix1* mutants, consistent with the loss-of-function mutation, but was restored to WT levels in the rescue group ($p=0.045$). Mitochondrial marker ATP5s levels showed no significant differences across groups, indicating that mitochondrial content was unaffected by the loss of *heix1* (Figure 1).





Figure 1. Western blot analysis of P-ERK, HEIX1, ERK, and ATP5s expression in S2 cells. (A) Elevated P-ERK levels in *heix1/heix1* mutants were normalized in the rescue group, highlighting *heix1*'s role in P-ERK regulation. (B) ATP5s levels remained stable across all groups, while P-ERK levels were markedly increased in *heix1/heix1* mutants, indicating specific activation of the P-ERK pathway. Total ERK levels were consistent across conditions.

The results highlight the impact of *heix1* mutations on apoptotic markers and ERK signaling pathways. Western blot analysis (panel A) shows elevated levels of active caspase 9 and phosphorylated ERK (P-ERK) in *heix1/heix1* mutants compared to wild-type (WT) controls (active caspase 9: 2.5-fold increase, $p=0.003$; P-ERK: 2.8-fold increase, $p=0.002$). These elevations indicate enhanced apoptosis and stress signaling in mutants. Total ERK and pro-caspase 9 levels remained stable across groups, confirming specific activation of apoptotic and stress pathways rather than changes in overall protein expression. The rescue group (**heix1/Df*; UAS, Gal4, *tubp>HEIX1*) exhibited normalized levels of active caspase 9 and P-ERK, similar to WT, indicating that *heix1* restoration mitigates apoptosis and stress signaling. Panel B provides immunofluorescent evidence of caspase 9 activation in *heix1/heix1* mutant and WT tissues. DAPI staining (blue) highlights nuclei, while caspase 9 activity (red) indicates apoptotic regions. The merged images demonstrate significantly increased colocalization of caspase 9 signals in mutant tissues compared to WT, corroborating the Western blot findings of heightened apoptotic activity in mutants (Figure 2).



disrupted cristae, and an irregular nuclear envelope, highlighting the impact of *heix1* mutations on cellular ultrastructure and homeostasis. Scale bars represent 500 nm.

The fluorescence microscopy analysis highlights apoptotic and oxidative stress markers in the lymph gland and imaginal disc tissues of *Drosophila*. In the lymph gland (top row), DAPI staining (blue) identifies nuclei, caspase activity (red) marks apoptotic regions, and the merged image confirms widespread apoptosis, as indicated by the colocalization of nuclear and caspase signals. In the imaginal disc (bottom row), DAPI staining visualizes nuclei, while CM-H2DCFDA (green) detects reactive oxygen species (ROS), signifying oxidative stress. The merged image illustrates ROS localization within the nuclear regions. These findings demonstrate elevated apoptosis and oxidative stress in mutant tissues, further implicating the role of *heix1* in maintaining cellular homeostasis (Figure 4).




Figure 4. Immunofluorescent analysis of apoptosis and oxidative stress in lymph gland and imaginal disc tissues of *Drosophila*. * (Top row) Lymph gland tissue shows DAPI-stained nuclei (blue), caspase-mediated apoptosis (red), and colocalization of both signals in the merged image. (Bottom row) Imaginal disc tissue highlights ROS production (green) detected by CM-H2DCFDA staining, with nuclear localization observed in the merged image. Scale bars represent 20 μ m.

The Western blot analysis highlights changes in signaling and stress response pathways in wild-type (WT), *heix1/heix1* mutants, and rescue groups (**heix1/Df*; UAS, Gal4, *tubp>HEIX1*). On the left panel, *heix1/heix1* mutants exhibit significantly reduced HEIX1 expression, confirming the loss-of-function mutation. These mutants also show elevated levels of phosphorylated JNK (P-JNK) and phosphorylated ERK (P-ERK) compared to WT (P-JNK: 2.3-fold increase, $p=0.003$; P-ERK: 2.8-fold increase, $p=0.001$), indicating hyperactivation of stress response pathways. Total ERK levels remain unchanged across groups, demonstrating that the observed activation is specific to P-JNK and P-ERK. The right panel reveals stable ATP5s levels across all groups, indicating no significant changes in mitochondrial content. ATF6, a marker of ER stress, is significantly elevated in *heix1/heix1* mutants compared to WT (2.5-fold increase, $p=0.002$), reflecting heightened unfolded protein response (UPR) activation. HEIX1 expression and stress marker levels in the rescue group are restored to near-WT levels (P-JNK: 1.2-fold WT,

p=0.045; P-ERK: 1.3-fold WT, p=0.048; ATF6: 1.2-fold WT, p=0.047). GADHF serves as the loading control, confirming consistent protein loading across all samples (Figure 5).




Figure 5. Western blot analysis of signaling and stress response proteins in wild-type, heix1/heix1 mutants, and rescue S2 cells. Elevated P-JNK, P-ERK, and ATF6 levels in heix1/heix1 mutants highlight hyperactivation of stress pathways and ER stress, while stable ATP5s levels confirm no mitochondrial abundance changes. Restoration of heix1 in the rescue group reduces stress markers, emphasizing its role in regulating cellular homeostasis. GADHF is used as a loading control.

Discussion

The most significant finding of this study is that heix1 loss-of-function mutations lead to hyperactivation of the P-ERK signaling pathway, resulting in increased apoptosis, ER stress, and mitochondrial dysfunction. Elevated levels of phosphorylated ERK (P-ERK), phosphorylated JNK (P-JNK), and apoptotic markers such as active caspase 9 in heix1/heix1 mutants indicate that heix1 plays a critical role in regulating cellular stress responses and maintaining homeostasis. Restoration of heix1 expression in the rescue group mitigated these effects, normalizing P-ERK and apoptotic signaling and underscoring the protective role of heix1 in cellular adaptation to stress.

These results are consistent with studies that have identified the P-ERK branch of the unfolded protein response (UPR) as a key regulator of cellular homeostasis. Kumar et al. demonstrated the importance of ER stress sensors such as IRE1, P-ERK, and ATF6 in coordinating ER-mitochondrial signaling and adaptation during stress (17). Similarly, Ong et al. reported that chronic ER stress triggers PERK activation through IRE1 signaling, linking prolonged stress to apoptosis, a mechanism that aligns with the elevated

P-ERK and ATF6 levels observed in heix1 mutants (19).

The elevated expression of CHOP, a pro-apoptotic transcription factor regulated by ATF4 during prolonged ER stress, was also observed in heix1/heix1 mutants. Rios-Fuller et al. emphasized the role of CHOP in apoptosis, highlighting how its upregulation leads to protein synthesis imbalance, oxidative stress, and eventual cell death (20, 21). These findings correlate with the mitochondrial swelling, fragmented cristae, and increased reactive oxygen species (ROS) production observed in heix1 mutants. Furthermore, Balakireva et al. and Alam et al. demonstrated the dual role of ER stress-mediated pathways in apoptosis and autophagy, further supporting the notion that heix1 mutations disrupt the balance between these processes (22, 23).

In addition to apoptosis, heix1/heix1 mutants exhibited morphological abnormalities such as melanotic tumors, supporting previous findings by Xia et al., who identified heix1 as a melanotic tumor suppressor

gene in *Drosophila* (24). These abnormalities align with mitochondrial dysfunction, as described by Casanova et al., who emphasized the critical role of mitochondria in regulating energy supply, apoptosis, and cellular differentiation (25). Our findings highlight how *heix1* mutations compromise mitochondrial integrity, contributing to cellular and organismal dysfunction.

The role of P-ERK signaling in lifespan regulation is also noteworthy. Balakireva et al. linked chronic P-ERK activation to reduced lifespan in *Drosophila*, findings consistent with the shortened lifespan observed in *heix1/heix1* mutants in this study (22). Chronic stress can be responsible for this lifetime ban by accumulated prototoxicity and continuous UPR activation. Interestingly, the rescue group reduced the P-ERC activity and generalized apoptotic signaling, which can contribute to improve cellular feasibility and extended survival. These findings emphasize protected genetic routes between *Drosophila* and mammals, which makes this model the organism ER stress and tumorigenesis to study. Munnik et al. The *Drosophila Melanogaster* was highlighted as a platform for the anticancer drug discovery, cited (26), citing high genetic similarity for its simple routes and humans. The results of this study strengthen this perception, as the regulatory role of the *heix1* in P-ERK and JNK routes provides valuable insight into the cellular stress system and their medical capacity. Finally, this study indicates that the *heix1* is necessary to regulate ER and mitochondrial homeostasis, reduce apoptosis, and prevent oxidative stress. The disadvantage of the *heix1* leads to ER stress, mitochondrial dysfunction and hyperactivation of apoptotic routes, as P-ERK, CHOP, and active caspase show increase in 9 levels. These results install the *Heix1* as an important regulator of cellular stress reactions and provide a foundation for the forward discovery of its role in ER stress, apoptosis and disease models associated with oxidative damage.

Conflict of interests: Authors declare that there is no possible struggle of interests related to research, writing or publication of this article.

Funding: Authors declare that no financial assistance was received for research, writing or publication of this article.

Acknowledgment

We are deeply grateful to God for His unfailing love, granting me health, knowledge, and patience throughout this journey. This work is dedicated to the honorable Prophet Mohammad, His household, and Imam Al-Mahdi (peace be upon him). We sincerely thank Dr. G.M. Rubin and the Berkeley *Drosophila* Genome Project, along with the Bloomington Stock Center, Patrik Verstreken, Tsinghua Fly Center, the Developmental Studies Hybridoma Bank, and Zhaohui Wang's lab for their support and resources. Special thanks to Zhengxing Wu for providing essential microscopy tools and facilities. We are indebted to my advisor, Professor Hong Ling, for his guidance and encouragement, and to my lab colleagues for their invaluable help in a new environment. We also acknowledge Misan University and the Republic of Iraq for financial support, as well as my family for their prayers, love, and unwavering support, which have been my foundation throughout this journey. This work is humbly dedicated to my fellow Iraqis and researchers worldwide.

References

- 1.Strunov A, Boldyreva LV, Andreyeva EN, Pavlova GA, Popova JV, Razuvaeva AV, et al. Ultrastructural analysis of mitotic Drosophila S2 cells identifies distinctive microtubule and intracellular membrane behaviors. BMC Biol. 2018;16(1):68.
- 2.Flatt T. Life-History Evolution and the Genetics of Fitness Components in Drosophila melanogaster. Genetics. 2020;214(1):3-48.
- 3.Su TT. Drug screening in Drosophila; why, when, and when not?. Wiley Interdiscip Rev Dev Biol. 2019;8(6):e346.
- 4.Matthew J, Vishwakarma V, Le TP, Agsunod RA, Chung S. Coordination of cell cycle and morphogenesis during organ formation. Elife. 2024;13:e95830.
- 5.Steinmetz EL, Noh S, Klöppel C, Fuhr MF, Bach N, Raffael ME, et al. Generation of Mutants from the 57B Region of Drosophila melanogaster. Genes (Basel). 2023;14(11):2047.
- 6.Yang Y, Lu D, Wang M, Liu G, Feng Y, Ren Y, et al. Endoplasmic reticulum stress and the unfolded protein response: emerging regulators in progression of traumatic brain injury. Cell Death Dis. 2024;15(2):156.
- 7.Rühmkorf A, Harbauer AB. Role of Mitochondria-ER Contact Sites in Mitophagy. Biomolecules. 2023;13(8):1198.
- 8.Nieblas B, Pérez-Treviño P, García N. Role of mitochondria-associated endoplasmic reticulum membranes in insulin sensitivity, energy metabolism, and contraction of skeletal muscle. Front Mol Biosci. 2022;9:959844.
- 9.Mainali N, Ayyadevara S, Ganne A, Shmookler Reis RJ, Mehta JL. Protein homeostasis in the aged and diseased heart. J Cardiovasc Aging. 2023;3(2):16.
- 10.Ottens F, Franz A, Hoppe T. Build-UPS and break-downs: metabolism impacts on proteostasis and aging. Cell Death Differ. 2021 Feb;28(2):505-521. doi: 10.1038/s41418-020-00682-y. Epub 2021 Jan 4. Erratum in: Cell Death Differ. 2022;29(2):465.
- 11.Chen X, Shi C, He M, Xiong S, Xia X. Endoplasmic reticulum stress: molecular mechanism and therapeutic targets. Signal Transduct Target Ther. 2023;8(1):352.
- 12.Smedley GD, Walker KE, Yuan SH. The Role of PERK in Understanding Development of Neurodegenerative Diseases. Int J Mol Sci. 2021;22(15):8146.
- 13.Jiang SY, Tang JJ, Xiao X, Qi W, Wu S, Jiang C, et al. Schnyder corneal dystrophy-associated UBIAD1 mutations cause corneal cholesterol accumulation by stabilizing HMG-CoA reductase. PLoS Genet. 2019;15(7):e1008289.
- 14.Jo Y, Hamilton JS, Hwang S, Garland K, Smith GA, Su S, et al. Schnyder corneal dystrophy-associated UBIAD1 inhibits ER-associated degradation of HMG CoA reductase in mice. Elife. 2019;8:e44396.
- 15.Li Y, Liu S, Wang YT, Min H, Adi D, Li XM, et al. TBL2 methylation is associated with hyper-low-density lipoprotein cholesterolemia: a case-control study. Lipids Health Dis. 2020;19(1):186.
- 16.Xu Z, Duan F, Lu H, Abdulkadhim Dragh M, Xia Y, Liang H, et al. UBIAD1 suppresses the proliferation of bladder carcinoma cells by regulating H-Ras intracellular trafficking via interaction with the C-terminal domain of H-Ras. Cell Death Dis. 2018;9(12):1170.
- 17.Kumar V, Maity S. ER Stress-Sensor Proteins and ER-Mitochondrial Crosstalk-Signaling Beyond (ER) Stress Response. Biomolecules. 2021;11(2):173.
- 18.Abdulkadhim Dragh M, Sabeeh Al-Allak Z, Zamil Gataa Allami Z. Cloning and expression of UbiA human gene using innovative methodologies for recombinant protein production in PUASt vector. Immunopathol Persa. 2025;11(1):e40643.

- 19.Ong G, Ragetli R, Mnich K, Doble BW, Kammouni W, Logue SE. IRE1 signaling increases PERK expression during chronic ER stress. *Cell Death Dis.* 2024;15(4):276.
- 20.Rios-Fuller TJ, Mahe M, Walters B, Abbadi D, Pérez-Baos S, Gadi A, et al. Translation Regulation by eIF2 α Phosphorylation and mTORC1 Signaling Pathways in Non-Communicable Diseases (NCDs). *Int J Mol Sci.* 2020;21(15):5301.
- 21.Dragh MA, Xu Z, Al-Allak ZS, Hong L. Vitamin K2 Prevents Lymphoma in *Drosophila*. *Sci Rep.* 2017;7(1):17047.
- 22.Balakireva Y, Nikitina M, Makhnovskii P, Kukushkina I, Kuzmin I, Kim A, et al. The Lifespan of *D. melanogaster* Depends on the Function of the Gagr Gene, a Domesticated gag Gene of *Drosophila* LTR Retrotransposons. *Insects.* 2024;15(1):68.
- 23.Alam R, Kabir MF, Kim HR, Chae HJ. Canonical and Noncanonical ER Stress-Mediated Autophagy Is a Bite the Bullet in View of Cancer Therapy. *Cells.* 2022;11(23):3773.
- 24.Xia Y, Midoun SZ, Xu Z, Hong L. Heixuedian (heix), a potential melanotic tumor suppressor gene, exhibits specific spatial and temporal expression pattern during *Drosophila* hematopoiesis. *Dev Biol.* 2015;398(2):218-30.
- 25.Casanova A, Wevers A, Navarro-Ledesma S, Pruimboom L. Mitochondria: It is all about energy. *Front Physiol.* 2023;14:1114231.
- 26.Munnik C, Xaba MP, Malindisa ST, Russell BL, Sooklal SA. *Drosophila melanogaster*: A platform for anticancer drug discovery and personalized therapies. *Front Genet.* 2022;13:949241.