



Thyroid Disorders Among People with Type 2 Diabetes in Misan Province: Prevalence Study

Rasha K Khudhur^{*1}, Ishraq J. Hasan², Maysaa Ghazi³, Malik M. Hasan²

¹Department of Pharmacology, Misan University, Faculty of Medicine, Misan, Iraq

²Department of Physiology, Misan University, Faculty of Medicine, Misan, Iraq

³Department of Microbiology, Misan University, Faculty of Medicine, Misan, Iraq



Article History:

Received on: 10 Jul 2020
Revised on: 11 Aug 2020
Accepted on: 19 Aug 2020

Keywords:

Diabetes mellitus,
hypothyroidism,
hyperthyroidism

ABSTRACT

Diabetes mellitus type 2 (DMT2) with thyroid disorders are common endocrine disorders, and both of them mostly come close in any clinical practice. Hormones of the thyroid gland can influence carbohydrate metabolism. On the other side, there is a degree of insulin resistance in DMT2. The study aimed is to determine the prevalence of thyroid dysfunctions in DMT2. A cross-sectional study was done enrolling 100 participants included, (n= 40) men, and (n= 60) women, their age ranged between 20-90 years old. Evaluation for thyroid status and assessment of T3, T4, and TSH levels were done. The results were 60% euthyroid, 33% hypothyroid, and 7% hyperthyroid cases from 100 patients with diabetes mellitus; among them, 17% in young adults, 49% in middle age, and 34% in the old age patients, the diabetic patients' results comprised between both sexes were 40% male and 60% females. In a comparison between rural and urban patients, the results were 44% rural and 56% urban. The study found there's 22% of thyroid dysfunction patients were with family history, and 42% of DM patients with family history, besides, there is 28% of patients suffered from hypertension and 33% of patients were smokers. So, the study showed a high prevalence of dysfunction among those with DMT2, especially hypothyroidism. Family history and DM might be pre-existing factors to the development of thyroid dysfunction. Hypertension and tobacco smoking might be a risk factor for thyroid dysfunction.

*Corresponding Author

Name: Rasha K Khudhur

Phone: 07710792262

Email: Medicalresearch77@yahoo.com

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v12i1.3939>

Production and Hosted by

Pharmascope.org

© 2021 | All rights reserved.

INTRODUCTION

The two most common endocrinopathies encountered in clinical practice are diabetes mellitus (DM)

and thyroid dysfunction (TD) (Sarode *et al.*, 2017), and the relation between them was first published in 1979 (Vikhe *et al.*, 2013). Thyroid dysfunction manifest as hyperthyroidism or hypothyroidism (Ahmed *et al.*, 2017). The metabolic process of normal growth and development regulated by thyroid hormones (Silva *et al.*, 2017). DM is accompanied by chronic hyperglycemia and associated with carbohydrate, lipid, and protein metabolism disturbances (Ahmed *et al.*, 2017). The prevalence of TD in diabetic patients is significantly higher than in healthy people (Manjunath *et al.*, 2013). There are many records that, show that iodothyronines are insulin antagonist with elevated levels being diabetogenic while absence might inhibit the development of DM (Uppal *et al.*, 2013).

MATERIALS AND METHODS

Study design and setting

A cross-sectional observational study conducted to determine the prevalence of TD in adult patients (aged 20-90 years) with DMT2, attending the outpatient's clinic at the Diabetes and Endocrinology Center at Misan, during 2018-2019.

Participants

A total of 100 DM participants, included, 40 men and 60 women. A complete history was taken, and clinical examination and laboratory investigations were done.

Inclusion criteria

1. Individuals with DMT2 were treated with oral hypoglycemic drugs
2. Those received insulin in addition to oral therapy.

Exclusion criteria

1. Those refused including in the study.
2. Uncontrol DM.
3. Uncomfortable patients.

Procedure

All patients were evaluated for thyroid status, and assessment of T3, T4 and TSH levels.

Normal values

1. T3 (0.9-2.3 nmol/l)
2. T4 (60-120 nmol/l)
3. TSH (0.25- 5.0 uIU/ml)

RESULTS

Of 100 DMT2 studied, 60 (60%) were found to have euthyroid, 7 (7%) hyperthyroid and the hypothyroid subjects were 33 (33%), presented in Table 1.

Table 2 showed the relation between the TD and age stages of DMT2 patients. The results showed that the younger age group was 17% (10% Euthyroid, 2% Hyperthyroid, and 5% Hypothyroid), whereas in the middle age, the results were 49% (30% Euthyroid, 3% Hyperthyroid, and 16% Hypothyroid). In the elderly, we found 34% (20% Euthyroid, 2% Hyperthyroid, 12% Hypothyroid).

In Table 3, we compared the female and male patients depending on the functional status of the thyroid gland, wherein male. It was: 26% Euthyroid,

3% Hyperthyroid, and 11% Hypothyroid. Whereas in female, it was 34% Euthyroid, 4% Hyperthyroid, and 22% Hypothyroid.

In Table 4, the results explained the distribution of cases based on the residential area where 44% in the rural areas (26% euthyroid, 3% hyperthyroid, 15% hypothyroid), and 56% in the urban regions (34% euthyroid, 4% hyperthyroid, 18% hypothyroid).

Table 5 shows the percentages of cases with a family history of thyroid disorders, 22% (5% Euthyroid, 3% Hyperthyroid, 14% Hypothyroid), while those with a family history were 42% (23% Euthyroid, 3% Hyperthyroid, 16% Hypothyroid).

Table 6 showed 28% of participants with hypertension (17% euthyroid, 2% hyperthyroid, 9% hypothyroid). 33% of cases were smoker (23% euthyroid, 2% hyperthyroid, 8% hypothyroid).

DISCUSSIONS

The TD had been reported to be highly prevalent in DMT2, and both hypothyroidism and hyperthyroidism are known to have adverse effects on glycaemic control (Petry, 2002). Our study entails screening for biochemical evidence of thyroid disorders in DMT2. The findings from the results were 40% thyroid dysfunctions (33% hypothyroid and 7% hyperthyroid), and this is consistent with previous similar studies performed by Telwani et al., Sarode et al., where their results were 29%, 31.2%, 32.4% respectively, in addition to the study of Sarode et al. found 29% patients were detected with thyroid disorders (22% hypothyroid and 7% hyperthyroid) (Telwani et al., 2017; Sarode et al., 2017). In Nigeria, there was a study had shown a high incidence reached to 46.5% (Udiong et al., 2007). The cause may be due to the fact of the DMT2 results from insulin resistance (Wang, 2013). Sometimes, the abnormality of thyroid hormones level associate with insulin resistance, lead to decline conversion of T4 to active T3, also decline hypothalamus thyrotropin-releasing hormone (TRH) in DM (Ahmed et al., 2017).

The results appeared that prevalence of DMT2 increased with older age group. These results agreed with studies of Gesing and Barbesino (Gesing, 2015; Barbesino, 2019). The ageing process decreases insulin sensitivity and alteration compensation of beta-cell function in the face of increasing insulin resistance (Chang and Halter, 2003). This decrement in beta cell proliferation capacity enhanced sensitivity to apoptosis (Maedler et al., 2006).

Noh et al. showed the initial and second phase of insulin secretion typically that which decrease at the

Table 1: Incidence of TD among DMT2.

TD	%
Euthyroid	60
Hyperthyroid	7
Hypothyroid	33

Table 2: Relation between the age and thyroid disorders in DMT2.

TD	Euthyroid	Hyperthyroid	Hypothyroid	Total
Age			%	
Young adult (18-35)	10	2	5	17
Middle age(36-55)	30	3	16	49
Old age> 55	20	2	12	34
Total	60	7	33	100

Table 3: Relation between sex and thyroid disorders in DMT2.

TD	Euthyroid	Hyperthyroid	Hypothyroid	Total
Sex			%	
Male	26	3	11	40
Female	34	4	22	60
Total	60	7	33	100

Table 4: Relation between the address and thyroid disorders in DMT2.

TD	Euthyroid	Hyperthyroid	Hypothyroid	Total
Address			%	
Rural	26	3	15	44
Urban	34	4	18	56
Total	60	7	33	100

Table 5: Relation between family history and thyroid disorders in DMT2.

TD	Euthyroid	Hyperthyroid	Hypothyroid	Total
Family history			%	
With thyroid dysfunction	5	3	14	22
With DMT2	23	3	16	42
Total	28	6	30	64

Table 6: Relation between the risk factors and TD in DMT2.

TD	Euthyroid	Hyperthyroid	Hypothyroid	Total
Risk factors			%	
Hypertension	17	2	9	28
smoking	23	2	8	33

rate of approximately 0.7% per year with the ageing process (Noh *et al.*, 2018).

In comparison between female and male depending on the functional status of the thyroid gland, the results were in male less than female. These results agreed with studies of Khurana *et al.* and Ding (Khurana *et al.*, 2016; Ding *et al.*, 2017). That woman have a higher incidence of TD than men (Meng *et al.*, 2015).

For the residential area, randomly selected cases of DMT2, which were 44% in the rural area and 56% in the urban area, the results were elevated of thyroid disorders in urban more than rural people. This result was agreed with Aung *et al.* (Aung *et al.*, 2018). The cause of DMT2 was higher in the more deprived areas relative to the more affluent areas (Connolly, 2000). In the study of Santos *et al.* described the socio-demographic variables, he found that the elderly individuals residing in the urban area displayed a more significant number of verified comorbidities. The old aged group DM is living in rural sites have better health (Santos *et al.*, 2013).

The study of Dudzińska *et al.* found differences between groups depending on the place of living, and this did not have a significant influence on the DM metabolic control. Moreover, these data are consistence with the previous study conducted by Childs (Dudzinska *et al.*, 2013; Childs, 2016).

The results appeared that there's 42% family history of DM, and this result agreed with many other studies such as Arslanian and Saad, Valdez and Liu (Arslanian *et al.*, 2005; Valdez *et al.*, 1999).

In case of risk factors, the results discovered 28% known facts of hypertension which correspond to other studies such as Saito and Saruta; and 33% smokers as a risk factor for both T2DM and TD, as the previous study of Chang, Maddatu *et al.*, they found that tobacco smoking, related with a variety disorder of endocrine systems (Chang, 2012; Maddatu *et al.*, 2017).

CONCLUSIONS

TD has a higher prevalence in DMT2, and more in women. Hypothyroidism was the most common disorder. Family history and DM might be pre-existing factors to develop thyroid dysfunction. Hypertension and tobacco smoking might be a risk factor for thyroid dysfunction.

Conflict of interest

The authors declare that they have no conflict of interest for this study.

Funding Support

The authors declare that they have no funding support for this study.

REFERENCES

- Ahmed, A. A., Mohamed, S. B., Elmadi, S. A., Abdorabo, A. A., Ismail, I. M., Ismail, A. M. 2017. Assessment of thyroid dysfunctions in type 2 diabetes mellitus patients in Surman, Western-Libya. *Int J Clin Exp Med Sci*, 3.
- Arslanian, S. A., Bacha, F., Saad, R., Gungor, N. 2005. Family History of Type 2 Diabetes Is Associated With Decreased Insulin Sensitivity and an Impaired Balance Between Insulin Sensitivity and Insulin Secretion in White Youth. *Diabetes Care*, 28(1):115–119.
- Aung, W. P., Htet, A. S., Bjertness, E., Stigum, H., Chongsuvivatwong, V., Kjøllesdal, M. K. R. 2018. Urban–rural differences in the prevalence of diabetes mellitus among 25–74 year-old adults of the Yangon Region, Myanmar: two cross-sectional studies. *BMJ Open*, 8(3):e020406–e020406.
- Barbesino, G. 2019. Thyroid Function Changes in the Elderly and Their Relationship to Cardiovascular Health: A Mini-Review. *Gerontology*, 65(1):1–8.
- Chang, A. M., Halter, J. B. 2003. Aging and insulin secretion. *American Journal of Physiology-Endocrinology and Metabolism*, 284(1):E7–E12.
- Chang, S. A. 2012. Smoking and type 2 diabetes mellitus. *Diabetes & metabolism journal*, 36(6):399–403.
- Childs, D. B. 2016. Comparison of Thyroid Disease Mortality between Urban and Rural Populations in Southwest Georgia. *Walden University*.
- Connolly, V. 2000. Diabetes prevalence and socioeconomic status: a population based study showing increased prevalence of type 2 diabetes mellitus in deprived areas. *Journal of Epidemiology & Community Health*, 54(3):173–177.
- Ding, X., Xu, Y., Wang, Y., Li, X., Lu, C., Su, J., Jin, Y 2017. Gender disparity in the relationship between the prevalence of thyroid nodules and metabolic syndrome components: the SHDC-CDPC community-based study. *Mediators of Inflammation*.
- Dudzinska, M., Tarach, J. S., Zwolak, A., Kurowska, M., Malicka, J., Smolen, A., Nowakowski, A. 2013. Type 2 diabetes mellitus concerning the place of residence: evaluation of selected aspects of socio-demographic status, course of diabetes and quality of life-a cross-sectional study. *Annals of Agricultural and Environmental Medicine*, 20(4):869–874.
- Gesing, A. 2015. The thyroid gland and the process of aging. *Thyroid Research*, 8(Suppl 1):A8–A8.

- Khurana, A., Dhoat, P., Jain, G. 2016. Prevalence of thyroid disorders in patients of type 2 diabetes mellitus. *J Indian Acad Clin Med*, 17(1):12-17.
- Maddatu, J., Anderson-Baucum, E., Evans-Molina, C. 2017. Smoking and the risk of type 2 diabetes. *Translational Research*, 184:101-107.
- Maedler, K., Schumann, D. M., Schulthess, F., Oberholzer, J., Bosco, D., Berney, T., Donath, M. Y. 2006. Ageing correlates with decreased β -cell proliferative capacity and enhanced sensitivity to apoptosis: a potential role for Fas and pancreatic duodenal homeobox-1. *Diabetes*, 55(9):2455-2462.
- Manjunath, S., Krishnamurthy, V., Puttaswamy, B., Prabhu, S., Vishwanathaiah, P. 2013. Prevalence of subclinical thyroid disorders in type 2 diabetes mellitus. *International Journal of Medicine and Public Health*, 3(4):330-330.
- Meng, Z., Liu, M., Zhang, Q., Liu, L., Song, K., Tan, J., Ren, X 2015. Gender and age impacts on the association between thyroid function and metabolic syndrome in. *Chinese. Medicine*, 94(50):2193-2193.
- Noh, J. W., Jung, J. H., Park, J. E., Lee, J. H., Sim, K. H., Park, J., Yoo, B, K. 2018. The relationship between age of onset and risk factors, including family history and lifestyle in Korean population with type 2 diabetes mellitus. *Journal of Physical Therapy Science*, 30(2):201-206.
- Petry, N. M. 2002. A Comparison of Young, Middle-Aged, and Older Adult Treatment-Seeking Pathological Gamblers. *The Gerontologist*, 42(1):92-99.
- Santos, E. A. D., Tavares, D. M. D. S., Rodrigues, L. R., Dias, F. A., Ferreira, P. C. D. S. 2013. Morbilidades y Calidad de Vida de ancianos con diabetes mellitus residentes en zonas rurales y urbanas. *Revista da Escola de Enfermagem da USP*, 47(2):393-400.
- Sarode, A., Patel, R., Pendse, M., Patil, S. 2017. Thyroid disorders in type 2 diabetes mellitus. *Med Pulse Int Med J*, 4:64-86.
- Silva, L. A. D., Wouk, J., Weber, V. M. R., Malfatti, C. R. M., Osiecki, R. 2017. The relation between diabetes mellitus, thyroid hormones and caffeine. *J. Appl. Pharm. Sci*, 7(03):212-216.
- Telwani, A. A., Wani, Z. H., Ashraf, Y., Shah, A. A. 2017. Prevalence of thyroid dysfunction in type 2 diabetes mellitus: a case control study. *International Journal of Research in Medical Sciences*, 5(10):4527-4527.
- Udiong, C. E. J., Udoh, A. E., Etukudoh, M. E. 2007. Evaluation of thyroid function in diabetes mellitus in Calabar, Nigeria. *Indian Journal of Clinical Biochemistry*, 22(2):74-78.
- Uppal, V., Vij, C., Bedi, G. K., Vij, A., Banerjee, B. D. 2013. Thyroid Disorders in Patients of Type 2 Diabetes Mellitus. *Indian Journal of Clinical Biochemistry*, 28(4):336-341.
- Valdez, R., Yoon, P. W., Liu, T., Khoury, M. J. 1999. Family history and prevalence of diabetes in the US population: the 6-year results from the National Health and Nutrition Examination Survey. *Diabetes care*, 30(10):2517-2522.
- Vikhe, V. B., Kanitkar, S. A., Tamakuwala, K. K., Gaikwad, A. N., Kalyan, M., Agarwal, R. R. 2013. Thyroid dysfunction in patients with type 2 diabetes mellitus at a tertiary care centre. *Natl J Med Res*, 3(4):377-80.
- Wang, C. 2013. The Relationship between Type 2 Diabetes Mellitus and Related Thyroid Diseases. *Journal of diabetes research*, 390534.